

AD-A053 293

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER
SINGULAR SEMI-LINEAR EQUATIONS IN $L_1(\mathbb{R})$. (U)
AUG 76 S D FISHER

F/G 12/1

DAA629-75-C-0024
NL

UNCLASSIFIED

MRC-TSR-1665

| OF |
AD
A053293



END
DATE
FILMED
6-78
DDC

AD A 053293

MRC Technical Summary Report #1665

SINGULAR SEMI-LINEAR EQUATIONS
IN $L^1(\mathbb{R})$

Stephen D. Fisher

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

August 1976

(Received May 7, 1976)

DDC FILE COPY

Form 1473

[Signature]

DDC
RECEIVED
APR 28 1978
Chr A

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

and

National Science Foundation
Washington, D. C.
20550

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

SINGULAR SEMI-LINEAR EQUATIONS IN $L^1(\mathbb{R})$

Stephen D. Fisher

Technical Summary Report #1665
August 1976

ABSTRACT

Let g be a positive continuous function on \mathbb{R} which tends to zero at $-\infty$ and which is not integrable over \mathbb{R} . The boundary-value problem $-u'' + g(u) = f$, $u'(\pm\infty) = 0$, is considered for $f \in L^1(\mathbb{R})$. We show that this problem can have a solution if and only if g is integrable at $-\infty$ and if this is so then the problem is solvable precisely when $\int_{-\infty}^{\infty} f(t)dt > 0$. Some extensions of this result are also given.

AMS (MOS) Subject Classification - 34B15

Key Words - Non-linear boundary-value problem

Work Unit Number 1 (Applied Analysis)

ACCESSION NO.	
NTIS	White Section <input checked="" type="checkbox"/>
DOC	Diff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
GEN	AVAIL. DOC. OR SPECIAL
A	

SINGULAR SEMI-LINEAR EQUATIONS IN $L^1(\mathbb{R})$

Stephen D. Fisher

In [2] M. G. Crandall and L. C. Evans show that the singular semi-linear problem

$$(*) \quad \begin{cases} -u''(x) + \beta(u(x)) = f(x), & -\infty < x < \infty \\ u'(\pm \infty) = 0 \\ u'' \in L^1(\mathbb{R}) \end{cases}$$

has a solution for each $f \in L^1(\mathbb{R})$ with $\int_{\mathbb{R}} f > 0$ if (and only if) β is integrable at $-\infty$. Here β is a given positive monotone increasing continuous function on \mathbb{R} . In fact, they discuss the more general situation when β is a maximal monotone graph. In this paper we consider several extensions of the problem (*) and provide another technique for proving that these equations have a solution. In particular, we recover the result of Crandall and Evans by different means.

Theorem 1. Let g be a positive continuous function on \mathbb{R} with

$$\lim_{t \rightarrow -\infty} g(t) = 0, \quad \int_{-\infty}^{\infty} g(s) ds \text{ divergent.}$$

Let $L_+^1 = \{f \in L^1(\mathbb{R}) : \int_{\mathbb{R}} f > 0\}$; for $f \in L_+^1$ consider the problem

$$(1) \quad \begin{cases} -u''(x) + g(u(x)) = f(x), & -\infty < x < \infty \\ u'' \in L^1(\mathbb{R}) \\ u'(\pm \infty) = 0. \end{cases}$$

The following are equivalent:

- (a) (1) has a solution for all $f \in L^1_+$
- (b) (1) has a solution for some $f \in L^1_+$
- (c) g is integrable at $-\infty$.

Proof. (a) implies (b) is trivial. To see that (b) implies (c) suppose there is a function u with $u'' \in L^1$, $u'(\pm \infty) = 0$, and

$$(2) \quad -u'' + g(u) = f$$

for some $f \in L^1$. Then $u' \in L^\infty$ and u tends to $-\infty$ at both $\pm \infty$ for

the following reason. Suppose there is a sequence $x_n \rightarrow \infty$ with

$\lim_n u(x_n) = L > -\infty$. Let $\{y_n\}$ be any other sequence of real numbers tending to $+\infty$. Then from (2) we get

$$-\frac{1}{2}(u'(y_n))^2 + \frac{1}{2}(u'(x_n))^2 + H(u(y_n)) - H(u(x_n)) = \int_{x_n}^{y_n} fu'$$

where

$$H(t) = \int_0^t g(s)ds.$$

Hence, $\lim_n H(u(y_n))$ exists and equals $H(u(L))$. Thus, $H(u(t))$ has a

limit at ∞ which implies that u has limit L at ∞ since H is strictly monotone. But then $g(u(t))$ tends to $g(L) > 0$ as $t \rightarrow \infty$ which contradicts the fact that $g(u(t))$ is in $L^1(\mathbb{R})$. An identical argument shows u tends to $-\infty$ at $-\infty$. With H as above we also have

$$\frac{1}{2} [(u'(y))^2 - (u'(0))^2] + H(u(0)) - H(u(y)) = \int_y^0 fu'$$

for each $y, y < 0$. Thus, $H(u(y))$ has a finite limit as $y \rightarrow -\infty$. Since $u(y) \rightarrow -\infty$ as $y \rightarrow -\infty$ we find that $H(s)$ has a finite limit as $s \rightarrow -\infty$ implying that g is integrable at $-\infty$. The proof that (c) implies (a) is the most difficult. The first step is to show that the set of those $f \in L^1_+$ for which (1) is solvable is closed in L^1_+ ; the second step is then obviously to show that the set of those $f \in L^1_+$ for which (1) is solvable is dense in L^1_+ . To prove the first assertion, let $f_n \rightarrow f$ in $L^1(\mathbb{R})$, with $f, f_n \in L^1_+$. Let u_n satisfy

$$(3a) \quad -u_n'' + g(u_n) = f_n$$

$$(3b) \quad u_n'(\pm\infty) = 0$$

$$(3c) \quad u_n'' \in L^1(\mathbb{R}).$$

Integrate both sides of (3a) from $-\infty$ to x and then from x to $+\infty$ and use the fact that $g \geq 0$. This gives $|u_n'(x)| \leq \|f\|_1 + 1$ for all large n and hence

$$(4) \quad \|u_n'\|_{L^\infty(\mathbb{R})} \leq A, \quad n = 1, 2, \dots$$

This in turn implies that $\{u_n\}$ is equicontinuous. We may assume, therefore, that $\{u_n\}$ converges uniformly on compact subsets of \mathbb{R} to either $+\infty$, or $-\infty$, or to a continuous function u . Set

$$G(x) = \int_{-\infty}^x g(s) ds.$$

For any $x \in \mathbb{R}$ and any n we have

$$\begin{aligned} G(u_n(x)) &= \int_{-\infty}^x g(u_n(t)) u_n'(t) dt \\ &= \frac{1}{2} [u_n'(x)]^2 + \int_{-\infty}^x f_n u_n' \\ &\leq A_1. \end{aligned}$$

Hence, $u_n(x) \leq C$ for all n and all x . Thus, it is obviously impossible that $\{u_n\}$ tends to $+\infty$. Suppose that $\{u_n\}$ tends to $-\infty$ uniformly on compact subsets of \mathbb{R} . Again we have

$$(5) \quad -\frac{1}{2} (u_n'(x))^2 + G(u_n(x)) = \int_{-\infty}^x f_n(t) u_n'(t) dt$$

and hence

$$(6) \quad 0 = \int_{-\infty}^{\infty} f_n u_n'.$$

We may assume that $\{u_n'\}$ converges weak-* in $L^\infty(\mathbb{R})$ to a function p and also that $\{u_n'(0)\}$ converges. Integrating (3a) from 0 to x we see that $u_n'(x)$ converges pointwise to $p(x)$ on \mathbb{R} . Hence, (5) and (6) yield

$$-\frac{1}{2} (p(x))^2 = \int_{-\infty}^x f p$$

and

$$0 = \int_{-\infty}^{\infty} fp.$$

Hence, p has a limit of 0 at both $+\infty$ and $-\infty$. Again from (3a) we obtain

$$u'_n(y) - u'_n(x) + \int_y^x g(u_n(t))dt = \int_y^x f_n(t)dt$$

so that

$$p(y) - p(x) = \int_y^x f(t)dt.$$

Now let $y \rightarrow -\infty$ and $x \rightarrow +\infty$; we find

$$0 < \int_{-\infty}^{\infty} f = p(-\infty) - p(+\infty) = 0,$$

a contradiction. Note that this argument is dependent on g in only a minor way. In particular, if $\{g_n\}$ is a sequence of positive continuous functions converging uniformly on compact subsets to a positive continuous function g which tends to 0 at $-\infty$ and which lies in $L^1(-\infty, 0]$ but not in $L^1(\mathbb{R})$ and if, say, $\{g_n\}$ increases to g on $(-\infty, \infty)$, then the functions v_n which satisfy

$$-v''_n + g_n(v_n) = f, \quad v'_n(\pm\infty) = 0, \quad f \in L^1_+$$

are equicontinuous and uniformly bounded on compact subsets of \mathbb{R} . We shall make use of this later on.

Returning to the functions $\{f_n\}$ and $\{u_n\}$ we see that $\{u_n\}$ converges uniformly on compact subsets of \mathbb{R} to a continuous function u . We clearly have $u''_n \rightarrow u''$ in L^1_{loc} so that u satisfies

$$(7) \quad -u'' + g(u) = f \text{ on } \mathbb{R}.$$

Fatou's lemma implies $g(u)$ is in $L^1(\mathbb{R})$ and hence $u'' \in L^1(\mathbb{R})$; thus u' has limits at both $\pm\infty$ and u tends to $-\infty$ at both $\pm\infty$ as in the implication (b) implies (c). From (5) and (6) we get

$$-\frac{1}{2}(u'(x))^2 + G(u(x)) = \int_{-\infty}^x fu'$$

and

$$0 = \int_{-\infty}^{\infty} fu'.$$

Hence, u' tends to 0 at both $\pm\infty$, so that u is a solution of (1).

Note also that

$$\begin{aligned} \int_{-\infty}^{\infty} |u'' + f| &= \int_{-\infty}^{\infty} (u'' + f) = \int_{-\infty}^{\infty} f \\ &\leq \|f\|_1 \end{aligned}$$

and hence

$$(8) \quad \|u''\|_1 \leq 2\|f\|_1.$$

The second assertion, that there is a dense set of $f \in L^1_+$ for which (1) is solvable, will be proved in the following way. Let f be a continuous function on \mathbb{R} in L^1_+ with support in the interval $I = [a, b]$. We shall show (1) is solvable for this f . We assume temporarily that g is C^1 on \mathbb{R} .

We shall need the following Proposition.

Proposition. Let $a < b$ and let g be a positive C^1 function on \mathbb{R} which is integrable at $-\infty$ and bounded at $+\infty$; set

$$G(x) = \int_{-\infty}^x g(s) ds.$$

Then for each α, β the initial value problem

$$(9) \quad \begin{cases} -v''(x) + g(v(x)) = f(x), & a < x < b, \quad f \in L^2(a, b) \\ v(a) = \alpha, \quad v'(a) = \beta \end{cases}$$

has a unique solution. If $\alpha_n \rightarrow \alpha$ and $\beta_n \rightarrow \beta$ and if v_n is the solution of (9) for (α_n, β_n) , then v_n converges uniformly to the solution v of (9) for (α, β) . Finally, the family $\{v_{\alpha\beta}\}$ of solutions of (9) corresponding to the initial values $\{(\alpha, \beta) : -\infty < \alpha \leq \alpha_0, |\beta| \leq M\}$ is equicontinuous on $[a, b]$.

Proof. Once the equicontinuity is established the existence and uniqueness follow from standard results; see [1], Chapter 1. To obtain the equicontinuity assertion (from which the second assertion also follows), we multiply the top equation in (9) by v' and integrate to obtain

$$-\frac{1}{2} (v'(x))^2 + G(v(x)) + \frac{1}{2} \beta^2 - G(\alpha) = \int_a^x f v'$$

so that if x_0 is chosen with $|v'(x_0)| = \|v'\|_\infty$ we have

$$\begin{aligned} \|v'\|_\infty^2 &\leq \beta^2 + 2G(\alpha) + 2G(v(x_0)) + A \|v'\|_\infty \\ &\leq \beta^2 + 2G(\alpha) + 2G(\alpha + (b-a) \|v'\|_\infty) + A \|v'\|_\infty \\ &\leq \beta^2 + 2G(\alpha) + A_0 + A_1(\alpha + (b-a) \|v'\|_\infty) + A \|v'\|_\infty \end{aligned}$$

for some constants A_0, A_1 depending only on g . Hence, $\|v'\|_\infty$ is bounded for $|\beta| \leq M$ and $-\infty < \alpha \leq \alpha_0$.

Conclusion of proof of Theorem 1. Let f be a continuous function in L^1_+ with support in the interval (a, b) . We shall show that (1) is solvable for this f . First, on $(-\infty, a]$ we show that the equation

$$(10) \quad g(u(x)) = u''(x)$$

$$u(a) = c_1, \quad u'(-\infty) = 0$$

has a solution. Let v be the function with

$$v'(t) = (2G(t))^{-1/2}, \quad -\infty < t < c_1$$

$$v(c_1) = a$$

where

$$G(x) = \int_{-\infty}^x g(s) ds.$$

Then v is increasing and has range $(-\infty, a]$. Let u be the inverse of v on $(-\infty, a]$, $u(v(t)) = t$. Thus

$$u(a) = c_1$$

and

$$u'(x) = 1/v'(t) = (2G(t))^{1/2}$$

or

$$(11) \quad u'(x) = (2G(u(x)))^{1/2}.$$

If we differentiate both sides of (11) we see that u satisfies (10).

Similarly, there is a solution of

$$u''(x) = g(u(x)) \quad b < x < \infty$$

$$u(b) = c_2, \quad u'(\infty) = 0$$

which satisfies

$$u'(x) = -(2G(u(x)))^{1/2}, \quad b < x < \infty.$$

Hence, to finish the proof of the theorem we need only show that there is a solution v of the equation

$$(12) \quad -v'' + g(v) = f \quad \text{on } (a, b)$$

with

$$(13) \quad (a) \quad v'(a) = (2G(v(a)))^{1/2}$$

$$(b) \quad v'(b) = -(2G(v(b)))^{1/2}.$$

Let v_t be the solution of (12) with $v(a) = t$ and $v'(a) = (2G(t))^{1/2}$ assured by the Proposition. (We temporarily assume that g is bounded at $+\infty$ if, in fact, it is not.) Then

$$\begin{aligned} v_t'(b) &= v_t'(a) + \int_a^b v_t''(s) ds \\ &= (2G(t))^{1/2} + \int_a^b g(v_t(s)) ds - \rho \end{aligned}$$

where $\rho = \int_a^b f(t)dt > 0$. To show that t may be chosen with $v_t'(b) = -(2G(v_t(b)))^{1/2}$ we consider

$$l(t) = (2G(t))^{1/2} + (2G(v_t(b)))^{1/2} + \int_a^b g(v_t(s))ds - \rho.$$

The Proposition implies l is continuous. We have

$$l(t) \geq -\rho + (2G(t))^{1/2}.$$

Since G is unbounded, there are values of t with $l(t) > 0$. Next let $t \downarrow -\infty$; by the equicontinuity of the functions $\{v_t\}$ we must have $v_t \rightarrow -\infty$ uniformly on $[a, b]$ so that $l(t) \rightarrow -\rho < 0$; hence, there is a t_0 at which $l(t_0) = 0$, and thus (12) is solvable with the boundary conditions (13).

We have now shown that (1) is solvable for all $f \in L_+^1$ under the assumption

$$(14) \quad g \in C^1(\mathbb{R}) \cap L^\infty(\mathbb{R}), \quad g \notin L^1(\mathbb{R}).$$

If g is merely positive and continuous on \mathbb{R} with $g \in L^1(-\infty, 0)$, $g \notin L^1(\mathbb{R})$, then there is a sequence $\{g_n\}$ of positive functions satisfying (14) which converge uniformly on compact subsets of \mathbb{R} to g and which also increase to g on $(-\infty, \infty)$. The comments made earlier show that the solutions $\{u_n\}$ of (1) with g_n in place of g converge to a solution of (1) for g . This completes the proof of Theorem 1.

Remark. The condition $g \notin L^1(\mathbb{R})$ is necessary as well as sufficient in order that Theorem 1 be valid. For suppose $g \in L^\infty(\mathbb{R}) \cap L^1(\mathbb{R})$; then the

function G is bounded. If f is supported on $[-1, 1]$ and if (1) has a solution for f , then (13) must hold with u in place of v so that

$$0 = \sqrt{2G(u(-1))} + \sqrt{2G(u(1))} + \int_{-1}^1 g(u(s))ds - \int_{-1}^1 f(s)ds.$$

The first three terms of this expression are bounded, independent of u , and hence the integral of f over \mathbb{R} can not exceed some fixed number depending only on g .

Theorem 2. Let g be a positive continuous function on \mathbb{R} with

$$(15) \quad \lim_{t \rightarrow -\infty} g(t) = 0, \quad g \notin L^1(\mathbb{R}).$$

Let $B(x)$ be a positive absolutely continuous function on \mathbb{R} with $B' \in L^1(\mathbb{R})$ and B bounded away from zero. For $f \in L^1_+$ consider the equation

$$(16) \quad \begin{cases} -u''(x) + B(x)g(u(x)) = f(x), & -\infty < x < \infty \\ u'' \in L^1(\mathbb{R}) \\ u'(\pm\infty) = 0. \end{cases}$$

Then (16) has a solution for each $f \in L^1_+$ if and only if g is integrable at $-\infty$.

Proof. If (16) is solvable for some $f \in L^1_+$ with support in $[-1, 1]$ then $u' > 0$ on $(-\infty, -1]$ and $u' < 0$ on $[1, \infty)$. It now follows very much as in Theorem 1 that u tends to $-\infty$ at $\pm\infty$ and that g is integrable at $-\infty$.

To show the sufficiency of the condition that g be integrable at $-\infty$ we first show that the equations

$$(17) \quad \begin{cases} u''(x) = B(x)g(u(x)), & |x| \geq a > 0 \\ u(-a) = c_1, \quad u(a) = c_2 \\ u'(\pm\infty) = 0 \end{cases}$$

have a solution. As in the proof of Theorem 1, the solution u must be monotone increasing for $-\infty < x < -a$ and monotone decreasing on (a, ∞) ; we shall only consider the details for the case $-\infty < x < -a$, the other case being entirely similar. We wish to find a continuous function v with

$$(18) \quad \begin{aligned} v'(t) &= (2 \int_{-\infty}^t B(v(s))g(s)ds)^{-1/2}, \quad -\infty < t < c_1 \\ v(c_1) &= -a. \end{aligned}$$

If such a v exists, then the inverse function u of v will satisfy

$$\begin{aligned} u'(x) &= (2 \int_{-\infty}^x B(r)g(u(r))u'(r)dr)^{1/2} \\ u(-a) &= c_1 \end{aligned}$$

and hence u will satisfy (17). To see that (18) has a solution let b_1 and b_2 be positive numbers with $b_1 \leq B(s) \leq b_2$ for all s and let ξ_N be the function defined by

$$\xi_N(t) = (2 \int_{-N}^t g(s)ds)^{-1/2}, \quad -N \leq t \leq c_1.$$

Let $\Omega_N = \{w \in C(-N, c_1) : (2b_2)^{-1/2} \xi_N(t) \leq w(t) \leq (2b_1)^{-1/2} \xi_N(t) \text{ for all } t \in [-N, c_1]\}$ and let T map Ω_N into Ω_N by

$$(Tw)(x) = (2 \int_{-N}^x B(\tilde{w}(s))g(s)ds)^{-1/2}$$

where

$$\tilde{w}'(t) = w(t), \quad \tilde{w}(c_1) = -a.$$

Clearly $Tw \in \Omega_N$; if $\{w_n\}$ is a bounded sequence in Ω_N , then $\{\tilde{w}_n\}$ is equicontinuous and uniformly bounded. Thus, T is a compact mapping and so has a fixed point w_N which must satisfy

$$w_N(x) = (2 \int_{-N}^x B(\tilde{w}_N(s))g(s)ds)^{-1/2}, \quad -N \leq x \leq c_1.$$

The functions $\{\tilde{w}_N\}$ are equicontinuous and uniformly bounded on compact subsets of $(-\infty, c_1]$ and so a subsequence, again denoted by $\{\tilde{w}_N\}$, converges uniformly on compact subsets of $(-\infty, c_1]$ to a function \tilde{w}_0 . But we also see that

$$\int_{-N}^x B(\tilde{w}_N(s))g(s)ds \rightarrow \int_{-\infty}^x B(\tilde{w}_0(s))g(s)ds$$

uniformly on compact subsets of $(-\infty, c_1]$. Hence, $w_N \rightarrow w_0$ uniformly on compacta; setting $v = \tilde{w}_0$ we see that v satisfies (18).

The remainder of the proof of Theorem 2 is like that of Theorem 1; the condition that $B' \in L^1(\mathbb{R})$ is used to prove that the sequence $\{u_n\}$ can not go to $-\infty$.

Corollary 3. Let $a(x) \in L^1(\mathbb{R})$, $f \in L^1(\mathbb{R})$, and let g be a positive continuous function satisfying (15). Consider the equation

$$(19) \quad \begin{cases} (i) & -u''(x) + a(x)u'(x) + g(u(x)) = f(x), \quad -\infty < x < \infty \\ (ii) & u'' \in L^1(\mathbb{R}) \\ (iii) & u'(\pm \infty) = 0 \end{cases}$$

Let g be integrable at $-\infty$ and set $w(x) = \exp[-\int_0^x a(s)ds]$. A necessary and sufficient condition that (19) be solvable is that

$$(20) \quad \int_{\mathbb{R}} f(x)w(x)dx > 0.$$

If (21) is solvable for all $f \in L^1$ satisfying (20), then g is integrable at $-\infty$.

Proof. Let $x = H(y)$ where H is the inverse of the function I defined by

$$I'(x) = 1/w(x)$$

$$I(0) = 0.$$

Then both H and I are 1-1 monotone increasing functions mapping \mathbb{R} onto \mathbb{R} and the substitution $v(y) = u(H(y))$ reduces (19) to

$$(21) \quad \begin{cases} -v''(y) + (H'(y))^2 g(v(y)) = (H'(y))^2 f(H(y)) \\ v'' \in L^1, \quad v'(\pm \infty) = 0 \end{cases}$$

which has a solution according to Theorem 2 precisely when

$$\begin{aligned} 0 &< \int_{-\infty}^{\infty} (H'(y))^2 f(H(y)) dy \\ &= \int_{-\infty}^{\infty} f(x)w(x)dx. \end{aligned}$$

Remark. Let β be a maximal monotone graph lying in the upper half-plane; that is, $\beta(x)$ is a subset of $\{y > 0\}$ for each $x \in \mathbb{R}$. Let $\beta^0(x) = \min\{y : y \in \beta(x)\}$. The result of Crandall and Evans is that if

$$\int_{-\infty}^a \beta^0(x) dx < \infty$$

for some $a \in D(\beta)$, then the equation

$$(22) \quad -u''(x) + \beta(u(x)) \ni f(x), \quad u'(\pm\infty) = 0, \quad f \in L^1_+$$

is solvable. This result also follows from Theorem 1 in the following way.

Let $\{\beta_n\}$ be a sequence of positive continuous monotone increasing functions which increase to β^0 on $D(\beta)$ and which increase to $+\infty$ off $D(\beta)$. The solutions $\{u_n\}$ of (1) with β_n in place of g then decrease on \mathbb{R} to a solution u of (22).

A final result related to Theorem 1 is presented below.

Theorem 4. Let g be a positive continuous function on \mathbb{R} satisfying

(15). For $f \in L^1(\mathbb{R})$ consider the equation

$$(23) \quad \begin{cases} u''(x) + g(u(x)) = f(x), & -\infty < x < \infty \\ u'' \in L^1(\mathbb{R}) \\ u'(-\infty) = \xi_1, \quad u'(+\infty) = \xi_2 \end{cases}$$

where

$$(24) \quad \int_{-\infty}^{\infty} f(x) dx = \rho > \xi_2 - \xi_1.$$

(a) Suppose g is integrable at $-\infty$. If (23) has a solution for some f with compact support (which necessarily satisfies (24)) then $\xi_1 > 0 > \xi_2$.

If (23) has a solution for $f \equiv 0$, then $\xi_1 = -\xi_2$.

(b) If g is integrable at $-\infty$ and if $\xi_1 > 0 > \xi_2$, then (23) has a solution for all f with

$$(25) \quad \xi_2 - \xi_1 < \rho = \int_{-\infty}^{\infty} f(x) dx \leq \min\{\xi_2, -\xi_1\}.$$

(c) If (23) has a solution for some f satisfying (24), then g is integrable at $-\infty$.

Proof. (a). If f has support in $[a, b]$, then $u''(x) < 0$ for $x < a$ and $x > b$. If $u'(-\infty) \leq 0$, then $u' < 0$ on $(-\infty, a)$ and hence u is decreasing on $(-\infty, a)$. However, u must tend to $-\infty$ at both $-\infty$ and $+\infty$ if u is a solution of (23) and thus u can not decrease on $(-\infty, a)$. Likewise, $u'(+\infty)$ must be negative. Further, if $u'' + g(u) \equiv 0$, then

$$(u'(x))^2 + 2G(u(x)) \equiv \text{const. on } (-\infty, \infty)$$

which clearly implies that $\xi_1 = -\xi_2$.

(c) is proved exactly as in Theorem 1.

(b) is the most difficult of the assertions. First, exactly as in Theorem 1, it can be shown that the set of those f satisfying (24) for which (23) is solvable is closed in $L^1(\mathbb{R})$. Next, we show that if f has compact support, say in (a, b) , and if f satisfies

$$(25)' \quad \xi_2 - \xi_1 < \rho = \int_{-\infty}^{\infty} f(x) dx < \min(-\xi_1, \xi_2)$$

then (23) has a solution. The key to this, as in Theorem 1, is to show two things: first that the equations

$$(26) \quad \begin{cases} u''(x) + g(u(x)) = 0, & x \notin [a, b] \\ u'(-\infty) = \xi_1, u'(\infty) = \xi_2 \end{cases}$$

have a solution which necessarily satisfies

$$(27) \quad \begin{aligned} u'(a) &= (\xi_1^2 - 2G(u(a)))^{1/2} \\ u'(b) &= -(\xi_2^2 - 2G(u(b)))^{1/2} \end{aligned}$$

and second that the equation

$$(28) \quad u''(x) + g(u(x)) = f(x), \quad a \leq x \leq b$$

is solvable subject to the non-linear boundary conditions (27). Both these assertions are proved as the similar statements are in the proof of Theorem 1.

Remark. The upper bound in (25) is not completely satisfactory; however, the situation for (23) is more involved than that of (1) as (a) shows.

The author would like to thank Prof. M. Crandall for a number of helpful comments on a preliminary version of this manuscript.

REFERENCES

- [1] E. A. Coddington and N. Levinson, Theory of Ordinary Differential Equations, McGraw-Hill, N. Y., 1955.
- [2] M. G. Crandall and L. C. Evans, A singular semilinear equation in $L^1(\mathbb{R})$, MRC Technical Summary Report #1566, Sept. 1975.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 MRC-TSR-1665	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9
4. TITLE (and Subtitle) 6 SINGULAR SEMI-LINEAR EQUATIONS IN $L^1(\mathbb{R})$.	5. TYPE OF REPORT & PERIOD COVERED Summary Report, no specific reporting period	
7. AUTHOR(s) 10 Stephen D. Fisher	6. PERFORMING ORG. REPORT NUMBER 15	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706 221 200	8. CONTRACT OR GRANT NUMBER DAAG29-75-C-0024 NSA-MPS 75-0550V	
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.	10. PROGRAM ELEMENT, PROJECT TASK AREA & WORK UNIT NUMBERS	12. REPORT DATE 11 August 1976
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 24p.	13. NUMBER OF PAGES 18	15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) * the integral from + infinity to - infinity of $f(t)dt > 0$		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 National Science Foundation Washington, D. C. 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Non-linear boundary-value problem - infinity + or - infinity as an element of $L^1(\mathbb{R})$		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let g be a positive continuous function on \mathbb{R} which tends to zero at $-\infty$ and which is not integrable over \mathbb{R} . The boundary-value problem $-u'' + g(u) = f$, $u'(\pm\infty) = 0$, is considered for $f \in L^1(\mathbb{R})$. We show that this problem can have a solution if and only if g is integrable at $-\infty$ and if this is so then the problem is solvable precisely when $\int_{-\infty}^{\infty} f(t)dt > 0$. Some extensions of this result are also given.		